

(19) World Intellectual Property Organization  
International Bureau




(43) International Publication Date  
12 January 2006 (12.01.2006)

PCT

(10) International Publication Number  
**WO 2006/004532 A1**

- (51) International Patent Classification<sup>7</sup>: **C07D 401/14**, A61K 31/506, A61P 11/00
- (21) International Application Number:  
PCT/SE2005/001092
- (22) International Filing Date: 4 July 2005 (04.07.2005)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
0401762-0 5 July 2004 (05.07.2004) SE
- (71) Applicant (for all designated States except US): **ASTRAZENECA AB** [SE/SE]; S-151 85 Södertälje (SE).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **GABOS, Balint** [SE/SE]; AstraZeneca R & D Lund, S-221 87 Lund (SE). **RIPA, Lena** [SE/SE]; AstraZeneca R & D Lund, S-221 87 Lund (SE). **STENVALL, Kristina** [SE/SE]; AstraZeneca R & D Lund, S-221 87 Lund (SE).
- (74) Agent: **ASTRAZENECA**; Global Intellectual Property, S-151 85 Södertälje (SE).
- (81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SI, SK, TR), OAPI (BI, BJ, CI, CG, CL, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:  
— with international search report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: **NOVEL HYDANTOIN DERIVATIVES FOR THE TREATMENT OF OBSTRUCTIVE AIRWAY DISEASES**



(57) Abstract: The invention provides compounds of formula (I), wherein R<sup>1</sup> and R<sup>2</sup> are as defined in the specification; processes for their preparation; pharmaceutical compositions containing them; a process for preparing the pharmaceutical compositions; and their use in therapy.

WO 2006/004532 A1

BEST AVAILABLE COPY

## Novel hydantoin derivatives for the treatment of obstructive airway diseases

The present invention relates to novel hydantoin derivatives, processes for their preparation, pharmaceutical compositions containing them and their use in therapy.

5

Metalloproteinases are a superfamily of proteinases (enzymes) whose numbers in recent years have increased dramatically. Based on structural and functional considerations these enzymes have been classified into families and subfamilies as described in N.M. Hooper (1994) FEBS Letters 354:1-6. Examples of metalloproteinases include the matrix metalloproteinases (MMPs) such as the collagenases (MMP1, MMP8, MMP13), the gelatinases (MMP2, MMP9), the stromelysins (MMP3, MMP10, MMP11), matrilysin (MMP7), metalloelastase (MMP12), enamelysin (MMP19), the MT-MMPs (MMP14, MMP15, MMP16, MMP17); the reprolysin or adamalysin or MDC family which includes the secretases and sheddases such as TNF converting enzymes (ADAM10 and TACE); the  
10   astacin family which include enzymes such as procollagen processing proteinase (PCP);  
15   and other metalloproteinases such as aggrecanase, the endothelin converting enzyme family and the angiotensin converting enzyme family.

Metalloproteinases are believed to be important in a plethora of physiological disease  
20   processes that involve tissue remodelling such as embryonic development, bone formation and uterine remodelling during menstruation. This is based on the ability of the metalloproteinases to cleave a broad range of matrix substrates such as collagen, proteoglycan and fibronectin. Metalloproteinases are also believed to be important in the processing, or secretion, of biological important cell mediators, such as tumour necrosis  
25   factor (TNF); and the post translational proteolysis processing, or shedding, of biologically important membrane proteins, such as the low affinity IgE receptor CD23 (for a more complete list see N. M. Hooper *et al.*, (1997) Biochem. J. 321:265-279).

Metalloproteinases have been associated with many diseases or conditions. Inhibition of  
30   the activity of one or more metalloproteinases may well be of benefit in these diseases or conditions, for example: various inflammatory and allergic diseases such as, inflammation of the joint (especially rheumatoid arthritis, osteoarthritis and gout), inflammation of the

gastro-intestinal tract (especially inflammatory bowel disease, ulcerative colitis and gastritis), inflammation of the skin (especially psoriasis, eczema, dermatitis); in tumour metastasis or invasion; in disease associated with uncontrolled degradation of the extracellular matrix such as osteoarthritis; in bone resorptive disease (such as osteoporosis and Paget's disease); in diseases associated with aberrant angiogenesis; the enhanced collagen remodelling associated with diabetes, periodontal disease (such as gingivitis), corneal ulceration, ulceration of the skin, post-operative conditions (such as colonic anastomosis) and dermal wound healing; demyelinating diseases of the central and peripheral nervous systems (such as multiple sclerosis); Alzheimer's disease; extracellular matrix remodelling observed in cardiovascular diseases such as restenosis and atherosclerosis; asthma; rhinitis; and chronic obstructive pulmonary diseases (COPD).

MMP12, also known as macrophage elastase or metalloelastase, was initially cloned in the mouse by Shapiro *et al* [1992, Journal of Biological Chemistry 267: 4664] and in man by the same group in 1995. MMP12 is preferentially expressed in activated macrophages, and has been shown to be secreted from alveolar macrophages from smokers [Shapiro *et al*, 1993, Journal of Biological Chemistry, 268: 23824] as well as in foam cells in atherosclerotic lesions [Matsumoto *et al*, 1998, Am. J. Pathol. 153: 109]. A mouse model of COPD is based on challenge of mice with cigarette smoke for six months, two cigarettes a day six days a week. Wild-type mice developed pulmonary emphysema after this treatment. When MMP12 knock-out mice were tested in this model they developed no significant emphysema, strongly indicating that MMP12 is a key enzyme in the COPD pathogenesis. The role of MMPs such as MMP12 in COPD (emphysema and bronchitis) is discussed in Anderson and Shinagawa, 1999, Current Opinion in Anti-inflammatory and Immunomodulatory Investigational Drugs 1(1): 29-38. It was recently discovered that smoking increases macrophage infiltration and macrophage-derived MMP-12 expression in human carotid artery plaques Kangavari [Matetzky S, Fishbein MC *et al*., Circulation 102:(18), 36-39 Suppl. S, Oct 31, 2000].

MMP9 (Gelatinase B; 92kDa TypeIV Collagenase; 92kDa Gelatinase) is a secreted protein which was first purified, then cloned and sequenced, in 1989 [S.M. Wilhelm *et al* (1989) J. Biol. Chem. 264 (29): 17213-17221; published erratum in J. Biol. Chem. (1990) 265

(36): 22570]. A recent review of MMP9 provides an excellent source for detailed information and references on this protease: T.H. Vu & Z. Werb (1998) (In : Matrix Metalloproteinases, 1998, edited by W.C. Parks & R.P. Mecham, pp. 115 – 148, Academic Press. ISBN 0-12-545090-7). The following points are drawn from that review  
5 by T.H. Vu & Z. Werb (1998).

The expression of MMP9 is restricted normally to a few cell types, including trophoblasts, osteoclasts, neutrophils and macrophages. However, the expression can be induced in these same cells and in other cell types by several mediators, including exposure of the  
10 cells to growth factors or cytokines. These are the same mediators often implicated in initiating an inflammatory response. As with other secreted MMPs, MMP9 is released as an inactive Pro-enzyme which is subsequently cleaved to form the enzymatically active enzyme. The proteases required for this activation *in vivo* are not known. The balance of active MMP9 versus inactive enzyme is further regulated *in vivo* by interaction with  
15 TIMP-1 (Tissue Inhibitor of Metalloproteinases -1), a naturally-occurring protein. TIMP-1 binds to the C-terminal region of MMP9, leading to inhibition of the catalytic domain of MMP9. The balance of induced expression of ProMMP9, cleavage of Pro- to active MMP9 and the presence of TIMP-1 combine to determine the amount of catalytically active MMP9 which is present at a local site. Proteolytically active MMP9 attacks substrates  
20 which include gelatin, elastin, and native Type IV and Type V collagens; it has no activity against native Type I collagen, proteoglycans or laminins.

There has been a growing body of data implicating roles for MMP9 in various physiological and pathological processes. Physiological roles include the invasion of  
25 embryonic trophoblasts through the uterine epithelium in the early stages of embryonic implantation; some role in the growth and development of bones; and migration of inflammatory cells from the vasculature into tissues.


MMP9 release, measured using enzyme immunoassay, was significantly enhanced in fluids  
30 and in AM supernatants from untreated asthmatics compared with those from other populations [Am. J. Resp. Cell & Mol. Biol., Nov 1997, 17 (5):583-591]. Also, increased MMP9 expression has been observed in certain other pathological conditions, thereby

implicating MMP9 in disease processes such as COPD, arthritis, tumour metastasis, Alzheimer's disease, multiple sclerosis, and plaque rupture in atherosclerosis leading to acute coronary conditions such as myocardial infarction.

- 5 A number of metalloproteinase inhibitors are known (see for example the reviews of MMP inhibitors by Beckett R.P. and Whittaker M., 1998, *Exp. Opin. Ther. Patents*, 8(3):259-282, and by Whittaker M. *et al*, 1999, *Chemical Reviews* 99(9):2735-2776).




WO 02/074767 discloses hydantoin derivatives of formula

10




that are useful as MMP inhibitors, particularly as potent MMP12 inhibitors. The following three compounds are specifically disclosed in WO 02/074767

15



We have now discovered a group of compounds that are inhibitors of metalloproteinases and are of particular interest in inhibiting MMPs such as MMP12 and MMP9. The compounds of the present invention have beneficial potency, selectivity and/or pharmacokinetic properties. The compounds of the present invention are within the generic scope of WO 02/074767 but are of a type not specifically exemplified therein.

In accordance with the present invention, there is therefore provided a compound of formula (I)



wherein

$R^1$  represents C1 to 2 alkyl, cyclopropyl,  $OCH_3$ ,  $SCH_3$  or  $OCF_3$ ; said alkyl or cyclopropyl group being optionally further substituted by one or more fluoro atoms; and

$R^2$  represents C1 to 3 alkyl;

and pharmaceutically acceptable salts thereof.

The compounds of formula (I) may exist in enantiomeric forms. It is to be understood that all enantiomers, diastereomers, racemates and mixtures thereof are included within the scope of the invention.

Compounds of formula (I) may also exist in various tautomeric forms. All possible tautomeric forms and mixtures thereof are included within the scope of the invention.

5 In one embodiment,  $R^1$  represents C1 to 2 alkyl or cyclopropyl; said alkyl or cyclopropyl group being optionally further substituted by one or more fluoro atoms.

In another embodiment,  $R^1$  represents C1 to 2 alkyl optionally further substituted by one or more fluoro atoms.

10

In one embodiment,  $R^1$  represents cyclopropyl optionally further substituted by one or more fluoro atoms.

In one embodiment,  $R^1$  represents cyclopropyl.

15

In one embodiment,  $R^1$  represents trifluoromethyl.

In one embodiment,  $R^1$  represents  $OCH_3$  or  $SCH_3$ .

20 In one embodiment,  $R^2$  represents methyl or ethyl. In one embodiment,  $R^2$  represents methyl.

In one embodiment,  $R^1$  represents C1 to 2 alkyl or cyclopropyl; said alkyl or cyclopropyl group being optionally further substituted by one or more fluoro atoms and  $R^2$  represents  
25 methyl or ethyl.

In one embodiment,  $R^1$  represents C1 to 2 alkyl or cyclopropyl; said alkyl or cyclopropyl group being optionally further substituted by one or more fluoro atoms and  $R^2$  represents methyl.

- 5 In one embodiment,  $R^1$  represents C1 to 2 alkyl optionally further substituted by one or more fluoro atoms and  $R^2$  represents methyl or ethyl.



In one embodiment,  $R^1$  represents  $CF_3$  and  $R^2$  represents methyl or ethyl.

- 10 In one embodiment,  $R^1$  represents cyclopropyl and  $R^2$  represents methyl or ethyl.

Unless otherwise indicated, the term "C1 to 3 alkyl" referred to herein denotes a straight or branched chain alkyl group having from 1 to 3 carbon atoms. Examples of such groups include methyl, ethyl, n-propyl and i-propyl. The term "C1 to 2 alkyl" denotes methyl or  
15 ethyl.

Examples of a C1 to 2 alkyl optionally further substituted by one or more fluoro atoms include  $CF_3$ ,  $CH_2F$ ,  $CH_2CF_3$ ,  $CF_2CH_3$  and  $CF_2CF_3$ .

- 20 Examples of a cyclopropyl ring optionally further substituted by one or more fluoro atoms include 1-fluoro-1-cyclopropyl, 2,2-difluoro-1-cyclopropyl and 2,3-difluoro-1-cyclopropyl:



25

Examples of compounds of the invention include:



- (5S)-5-({[4-[(2-cyclopropylpyrimidin-5-yl)ethynyl]-3,6-dihydropyridin-1(2H)-yl]sulfonyl}methyl)-5-methylimidazolidine-2,4-dione;  
 (5S)-5-methyl-5-({[4-{{[2-(methylthio)pyrimidin-5-yl]ethynyl}-3,6-dihydropyridin-1(2H)-yl]sulfonyl}methyl}imidazolidine-2,4-dione;  
 5 (5S)-5-methyl-5-({[4-{{[2-(trifluoromethyl)pyrimidin-5-yl]ethynyl}-3,6-dihydropyridin-1(2H)-yl]sulfonyl}methyl}imidazolidine-2,4-dione;  
 (5S)-5-methyl-5-({[4-[(2-methylpyrimidin-5-yl)ethynyl]-3,6-dihydropyridin-1(2H)-yl]sulfonyl}methyl)imidazolidine-2,4-dione;  
 (5S)-5-({[4-[(2-ethylpyrimidin-5-yl)ethynyl]-3,6-dihydropyridin-1(2H)-yl]sulfonyl}methyl)-5-methylimidazolidine-2,4-dione;  
 10 (5S)-5-({[4-[(2-methoxypyrimidin-5-yl)ethynyl]-3,6-dihydropyridin-1(2H)-yl]sulfonyl}methyl)-5-methylimidazolidine-2,4-dione;  
 and pharmaceutically acceptable salts thereof.


- 15 Each exemplified compound represents a particular and independent aspect of the invention.

The compounds of formula (I) may exist in enantiomeric forms. Therefore, all enantiomers, diastereomers, racemates and mixtures thereof are included within the scope of the invention.

- 20 The various optical isomers may be isolated by separation of a racemic mixture of the compounds using conventional techniques, for example, fractional crystallisation, or HPLC. Alternatively the optical isomers may be obtained by asymmetric synthesis, or by synthesis from optically active starting materials.

- 25 Where optically isomers exist in the compounds of the invention, we disclose all individual optically active forms and combinations of these as individual specific embodiments of the invention, as well as their corresponding racemates.

Preferably the compounds of formula (I) have (5S)-stereochemistry as shown below:



Where tautomers exist in the compounds of the invention, we disclose all individual tautomeric forms and combinations of these as individual specific embodiments of the invention.

The present invention includes compounds of formula (I) in the form of salts. Suitable salts include those formed with organic or inorganic acids or organic or inorganic bases. Such salts will normally be pharmaceutically acceptable salts although non-pharmaceutically acceptable salts may be of utility in the preparation and purification of particular compounds. Such salts include acid addition salts such as hydrochloride, hydrobromide, citrate, tosylate and maleate salts and salts formed with phosphoric acid or sulphuric acid. In another aspect suitable salts are base salts such as an alkali metal salt, for example, sodium or potassium, an alkaline earth metal salt, for example, calcium or magnesium, or an organic amine salt, for example, triethylamine.

Salts of compounds of formula (I) may be formed by reacting the free base or another salt thereof with one or more equivalents of an appropriate acid or base.

The compounds of formula (I) are useful because they possess pharmacological activity in animals and are thus potentially useful as pharmaceuticals. In particular, the compounds of the invention are metalloproteinase inhibitors and may thus be used in the treatment of diseases or conditions mediated by MMP12 and/or MMP9 such as asthma, rhinitis, chronic obstructive pulmonary diseases (COPD), arthritis (such as rheumatoid arthritis and osteoarthritis), atherosclerosis and restenosis, cancer, invasion and metastasis, diseases involving tissue destruction, loosening of hip joint replacements, periodontal disease,

fibrotic disease, infarction and heart disease, liver and renal fibrosis, endometriosis, diseases related to the weakening of the extracellular matrix, heart failure, aortic aneurysms, CNS related diseases such as Alzheimer's disease and Multiple Sclerosis (MS), and hematological disorders.

5

In general, the compounds of the present invention are potent inhibitors of MMP9 and MMP12. The compounds of the present invention also show good selectivity with respect to a relative lack of inhibition of various other MMPs such as MMP8, MMP14 and MMP19. In addition, the compounds of the present invention also, in general, have improved log D values, in particular, having log D values in the range of  $0.5 < \log D < 2.0$ . Log D is a parameter that reflects the lipophilicity of a compound at physiological pH. As a consequence of these favourable log D values, the compounds of the present invention possess improved solubility characteristics and reduced plasma protein binding, leading to improved pharmacokinetic and pharmacodynamic properties.

15

Accordingly, the present invention provides a compound of formula (I), or a pharmaceutically acceptable salt thereof, as hereinbefore defined for use in therapy.

In another aspect, the invention provides the use of a compound of formula (I), or a pharmaceutically acceptable salt thereof, as hereinbefore defined in the manufacture of a medicament for use in therapy.

In another aspect, the invention provides the use of a compound of formula (I), or a pharmaceutically acceptable salt thereof, as hereinbefore defined in the manufacture of a medicament for use in the treatment of diseases or conditions in which inhibition of MMP12 and/or MMP9 is beneficial.

In another aspect, the invention provides the use of a compound of formula (I), or a pharmaceutically acceptable salt thereof, as hereinbefore defined in the manufacture of a medicament for use in the treatment of inflammatory disease.

30

In another aspect, the invention provides the use of a compound of formula (I), or a pharmaceutically acceptable salt thereof, as hereinbefore defined in the manufacture of a medicament for use in the treatment of an obstructive airways disease such as asthma or COPD.

5

In the context of the present specification, the term "therapy" also includes "prophylaxis" unless there are specific indications to the contrary. The terms "therapeutic" and "therapeutically" should be construed accordingly.

- 10 Prophylaxis is expected to be particularly relevant to the treatment of persons who have suffered a previous episode of, or are otherwise considered to be at increased risk of, the disease or condition in question. Persons at risk of developing a particular disease or condition generally include those having a family history of the disease or condition, or those who have been identified by genetic testing or screening to be particularly
- 15 susceptible to developing the disease or condition.

The invention further provides a method of treating a disease or condition in which inhibition of MMP12 and/or MMP9 is beneficial which comprises administering to a patient a therapeutically effective amount of a compound of formula (I) or a

20 pharmaceutically acceptable salt thereof as hereinbefore defined.

The invention also provides a method of treating an obstructive airways disease, for example, asthma or COPD, which comprises administering to a patient a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt

25 thereof as hereinbefore defined.

For the above-mentioned therapeutic uses the dosage administered will, of course, vary with the compound employed, the mode of administration, the treatment desired and the disorder to be treated. The daily dosage of the compound of formula (I)/salt (active

30 ingredient) may be in the range from 0.001 mg/kg to 75 mg/kg, in particular from 0.5 mg/kg to 30 mg/kg. This daily dose may be given in divided doses as necessary.

Typically unit dosage forms will contain about 1 mg to 500 mg of a compound of this invention.

The compounds of formula (I) and pharmaceutically acceptable salts thereof may be used  
5 on their own but will generally be administered in the form of a pharmaceutical  
composition in which the formula (I) compound/salt (active ingredient) is in association  
with a pharmaceutically acceptable adjuvant, diluent or carrier. Depending on the mode of  
administration, the pharmaceutical composition will preferably comprise from 0.05 to 99  
%w (per cent by weight), more preferably from 0.10 to 70 %w, of active ingredient, and,  
10 from 1 to 99.95 %w, more preferably from 30 to 99.90 %w, of a pharmaceutically  
acceptable adjuvant, diluent or carrier, all percentages by weight being based on total  
composition. Conventional procedures for the selection and preparation of suitable  
pharmaceutical formulations are described in, for example, "Pharmaceuticals - The Science  
of Dosage Form Designs", M. E. Aulton, Churchill Livingstone, 1988.

15 Thus, the present invention also provides a pharmaceutical composition comprising a  
compound of formula (I) or a pharmaceutically acceptable salt thereof as hereinbefore  
defined in association with a pharmaceutically acceptable adjuvant, diluent or carrier.


20 The invention further provides a process for the preparation of a pharmaceutical  
composition of the invention which comprises mixing a compound of formula (I) or a  
pharmaceutically acceptable salt thereof as hereinbefore defined with a pharmaceutically  
acceptable adjuvant, diluent or carrier.

25 The pharmaceutical compositions of this invention may be administered in a standard  
manner for the disease or condition that it is desired to treat, for example by oral, topical,  
parenteral, buccal, nasal, vaginal or rectal administration or by inhalation. For these  
purposes the compounds of this invention may be formulated by means known in the art  
into the form of, for example, tablets, capsules, aqueous or oily solutions, suspensions,  
30 emulsions, creams, ointments, gels, nasal sprays, suppositories, finely divided powders or  
aerosols for inhalation, and for parenteral use (including intravenous, intramuscular or  
infusion) sterile aqueous or oily solutions or suspensions or sterile emulsions.

In addition to the compounds of the present invention the pharmaceutical composition of this invention may also contain, or be co-administered (simultaneously or sequentially) with, one or more pharmacological agents of value in treating one or more diseases or conditions referred to hereinabove such as "Symbicort" (trade mark) product.


The present invention further provides a process for the preparation of a compound of formula (I) or a pharmaceutically acceptable salt thereof as defined above which, comprises:

a) reaction of a compound of formula (II)



wherein  $R^2$  is as defined in formula (I) and  $L^1$  represents a leaving group, with a


compound of formula (III) (or a salt thereof)



wherein  $R^1$  is as defined in formula (I); or


b) reaction of a compound of formula (X)

14




wherein  $R^2$  is as defined in formula (I),  $R^3$  is H or a suitable protecting group and X is a leaving group such as halide or triflate; with an acetylenic compound of formula (IX)

5



wherein  $R^1$  is as defined in formula (I); or


c) reaction of a compound of formula (XI)



10

wherein R represents H or trimethylsilyl,  $R^2$  is as defined in formula (I) and  $R^3$  represents H or a suitable protecting group; with an aryl halide or triflate of formula (VI)

15



(VI)

wherein R<sup>1</sup> is as defined in formula (I) and X represents halide or triflate;  
and optionally thereafter forming a pharmaceutically acceptable salt thereof.


5

In the above process (a), suitable leaving groups L<sup>1</sup> include halo, particularly chloro. The reaction is preferably performed in a suitable solvent optionally in the presence of an added base for a suitable period of time, typically 0.5 to 24 h, at ambient to reflux temperature. Typically solvents such as pyridine, dimethylformamide, tetrahydrofuran, acetonitrile or dichloromethane are used. When used, the added base may be an organic base such as triethylamine, diisopropylethylamine, N-methylmorpholine or pyridine, or an inorganic base such as an alkali metal carbonate. The reaction is typically conducted at ambient temperature for 0.5 to 16 h, or until completion of the reaction has been achieved, as determined by chromatographic or spectroscopic methods. Reactions of sulfonyl halides with various primary and secondary amines are well known in the literature, and the variations of the conditions will be evident for those skilled in the art.

15

Sulfonylchlorides of formula (II) (wherein L<sup>1</sup> represents chlorine) are conveniently prepared by oxidative chlorination of compounds of formula (IV)

20




(IV)



using methods that will be readily apparent to those skilled in the art (Mosher, J., *J. Org. Chem.* **1958**, 23, 1257; Griffith, O., *J. Biol. Chem.* **1983**, 258, (3), 1591; WO 02/074767).

Compounds of formula (III) can be prepared by various methods described in the literature or variations thereon as will be appreciated by those skilled in the art of synthetic organic chemistry. Suitable methods include, but are not limited to, those described below and are shown in Scheme 1.



Scheme 1

In Scheme 1, PG represents a suitable protecting group such as t-Boc; X represents a leaving group such as a halide or a triflate; R represents hydrogen or trimethylsilyl; tms represents trimethylsilyl; Ar represents a 5-pyrimidinyl ring substituted at the 2-position by R<sup>1</sup>; and R<sup>1</sup> is as defined in formula (I).

The reaction between the aryl- or vinyl derivative [(V) or (VI)] and an acetylene [(VII), (VIII) or (IX)] can be accomplished, optionally in a suitable solvent, using a catalyst such as a suitable palladium salt, for example,  $\text{PdCl}_2(\text{PPh}_3)_2$ , with/or without an added copper salt and with an amine base such as piperidine, triethylamine, diisopropylamine or diisopropylethylamine. When used, the added solvent may be, for example, tetrahydrofuran, acetonitrile or N,N-dimethylformamide. The reaction is conducted at ambient to reflux temperature for 20 minutes to several hours until chromatographic or spectroscopic methods indicate completion of the reaction. Palladium catalysed reactions involving acetylenic compounds are well known in the literature, and variations of the conditions will be evident for those skilled in the art. General methodology of this type is described in, for example, Brandsma, L., *Synthesis of Acetylenes, Allenes and Cumulenes: Methods and Techniques*, 2004, Elsevier Academic Press, chapter 16, pages 293-317; *Transition Metals-Catalysed Couplings of Acetylenes with  $sp^2$ -halides*, Sonogashira, K., *J. Organomet. Chem.*, **2002**, 653, 46-49; Tykwinski, R. R., *Angew. Chem. Int.Ed.*, **2003**, 42, 1566-1568.

The vinyl triflate (V) wherein X is O-triflate and PG is t-Boc can be prepared as described in the literature (Wustrow, D. J., *Synthesis*, **1991**, 993-995).

Suitable substituted pyrimidinyl halides or triflates of formula (VI) can be prepared by various methods described in the literature, for example, Budesinsky, Z. et al., *Coll. Czech. Chem. Commun.*, **1949**, 14, 223-235; Takahashi et al., *Chem. Pharm. Bull.*, **1958**, 6, 334-337; US 4,558,039.

The acetylenic compound (VIII) can be prepared from the triflate (V) via a palladium catalysed coupling reaction with trimethylsilylacetylene followed by, if necessary, deprotection of the trimethylsilyl group using, for example, potassium fluoride in a suitable solvent. Alternatively, preparation of compound (VIII) wherein R is H and PG is t-Boc can be accomplished by dehydrating a compound of formula (VII), for example, by mesylation followed by treatment with a suitable base, for example, diisopropylethylamine.

Acetylenic heteroaryl compounds of formula (IX) can be prepared by various methods described in the literature.

In process (b), the reactions are carried out using methods similar to those described above  
5 for the preparation of compounds of formula (VIII). If necessary, one nitrogen in the  
hydantoin ring of compounds of formula (X) can be protected using SEMCl ( $R^3 = \text{SEM}$ )  
before the palladium catalysed reaction is performed. Compounds of formula (X) can be  
prepared by acid catalysed deprotection of compounds of formula (V) ( $\text{PG} = \text{t-Boc}$ ),  
followed by reaction with a compound of formula (II), in the same way as described above  
10 for the preparation of compounds of formula (I).

In process (c), the reactions are carried out in a similar manner to those described above for  
the preparation of compounds of formula (VIII). If necessary, one nitrogen of the  
hydantoin ring of compounds of formula (XI) can be protected using SEMCl ( $R^3 = \text{SEM}$ )  
15 before the palladium catalysed reaction is performed. Compound (XI) is conveniently  
prepared from compound (VIII) wherein R is trimethylsilyl and PG is t-Boc by acid  
catalysed removal of the t-Boc group (for example, using acetyl chloride in methanol),  
followed by reaction with a compound of formula (II), as described above for the reaction  
between compounds of formulae (II) and (III).

20 It will be appreciated by those skilled in the art that in the processes of the present  
invention certain potentially reactive functional groups such as hydroxyl or amino groups  
in the starting reagents or intermediate compounds may need to be protected by suitable  
protecting groups. Thus, the preparation of the compounds of the invention may involve, at  
25 various stages, the addition and removal of one or more protecting groups.

Suitable protecting groups and details of processes for adding and removing such groups  
are described in 'Protective Groups in Organic Chemistry', edited by J.W.F. McOmie,  
Plenum Press (1973) and 'Protective Groups in Organic Synthesis', 3rd edition, T.W.  
30 Greene and P.G.M. Wuts, Wiley-Interscience (1999).

The compounds of the invention and intermediates thereto may be isolated from their reaction mixtures and, if necessary further purified, by using standard techniques.

The present invention will now be further explained by reference to the following  
5 illustrative examples.

#### General Methods

<sup>1</sup>H NMR and <sup>13</sup>C NMR spectra were recorded on a Varian *Inova* 400 MHz or a Varian  
10 *Mercury-VX* 300 MHz instrument. The central peaks of chloroform-*d* ( $\delta_H$  7.27 ppm),  
dimethylsulfoxide-*d*<sub>6</sub> ( $\delta_H$  2.50 ppm), acetonitrile-*d*<sub>3</sub> ( $\delta_H$  1.95 ppm) or methanol-*d*<sub>4</sub> ( $\delta_H$  3.31  
ppm) were used as internal references. Column chromatography was carried out using  
silica gel (0.040-0.063 mm, Merck). A Kromasil KR-100-5-C<sub>18</sub> column (250 × 20 mm,  
Akzo Nobel) and mixtures of acetonitrile/water with 0.1 % TFA at a flow rate of 10  
15 mL/min were used for preparative HPLC. Unless stated otherwise, starting materials were  
commercially available. All solvents and commercial reagents were of laboratory grade  
and were used as received.

The following method was used for LC/MS analysis:

20 Instrument Agilent 1100; Column Waters Symmetry 2.1 × 30 mm; Mass APCI; Flow rate  
0.7 mL/min; Wavelength 254 or 220 nm; Solvent A: water + 0.1% TFA; Solvent B:  
acetonitrile + 0.1% TFA ; Gradient 15-95%/B 2.7 min, 95% B 0.3 min.

The following method was used for LC analysis:

25 Method A. Instrument Agilent 1100; Column: Kromasil C18 100 × 3 mm, 5μ particle size,  
Solvent A: 0.1%TFA/water, Solvent B: 0.08%TFA/acetonitrile Flow rate 1 mL/min,  
Gradient 10-100%/B 20 min, 100% B 1 min. Absorption was measured at 220, 254 and  
280 nm.

Method B. Instrument Agilent 1100; Column: XTerra C 8, 100 × 3 mm, 5μ particle size,  
30 Solvent A: 15mM NH<sub>3</sub>/water, Solvent B: acetonitrile Flow rate 1 mL/min, Gradient 10-  
100%/B 20 min, 100% B 1 min. Absorption was measured at 220, 254 and 280 nm.

## Abbreviations:

Ac acetyl

DMF *N,N*-dimethylformamide

DMSO dimethyl sulfoxide

5 eq. equivalent

Et ethyl

LDA lithium diisopropyl amide

Me methyl

MS mass spectroscopy

10 *tert* tertiary

THF tetrahydrofuran

TFA trifluoroacetic acid

Example 1

15

(5*S*)-5-({[4-[(2-Cyclopropylpyrimidin-5-yl)ethynyl]-3,6-dihydropyridin-1(2*H*)-yl]sulfonyl}methyl)-5-methylimidazolidine-2,4-dione

The title compound was prepared following the general method of Yamanaka et al, *Synth. Commun.*, 1983, 312-314. To 5-bromo-2-cyclopropylpyrimidine (110 mg, 0.55 mmol) and (5*S*)-5-{{[4-ethynyl-3,6-dihydropyridin-1(2*H*)-yl]sulfonyl}methyl}-5-methylimidazolidine-2,4-dione (180 mg, 0.61 mmol) in THF (3 mL) was added Et<sub>3</sub>N (1 mL) and DMF (1 mL) at 35 °C. After a solution was formed, CuI (4 mol %) and PdCl<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub> (2 mol %) were added and the mixture was heated at 72 °C for 6 hours. The mixture was partitioned between EtOAc (15 mL) and water (10 mL), and the aqueous layer was extracted three times with EtOAc. The combined organic layers were dried and concentrated to give the crude product as a yellow oil. The title compound (65 mg) was obtained by purification using preparative HPLC.

<sup>1</sup>H-NMR (DMSO-*d*<sub>6</sub>): δ 10.75 (1H, s); 8.72 (2H, s); 8.03 (1H, s); 6.28 (1H, m); 3.84 (2H, m); 3.47 (2H, q); 3.30 (2H, m); 2.37 (2H, m); 2.21 (1H, m); 1.33 (3H, s); 1.10 (2H, m); 1.02 (2H, m).

APCI-MS m/z: 416 [MH<sup>+</sup>].

a) 5-Bromo-2-cyclopropylpyrimidine

5-Bromo-2-cyclopropylpyrimidine was prepared by the method of Budesinsky, Z., *Coll.*

5 *Czech. Chem. Commun.*, **1949**, *14*, 223-235. Cyclopropanecarboximidamide hydrochloride (2.5 g, 20.7 mmol) was dissolved in EtOH (4 mL), freshly prepared 4.1M NaOEt in EtOH (4.8 mL) was added followed by mucobromic acid (2.7 g, 10.3 mmol). The mixture was heated to 56 °C for 30 minutes, more NaOEt in EtOH (4.1M, 3.2 mL) was added and the reaction was stirred at 56 °C for another 15 minutes and then at room temperature overnight. The solvent was evaporated off, aqueous HCl (2M, 10 mL) was added and the brown solid was filtered off. The aqueous layer was extracted three times with dichloromethane. The combined organic layers were dried and concentrated to give a brown oil that together with the solid gave crude intermediate 5-bromo-2-cyclopropylpyrimidine-4-carboxylic acid (1.6 g). The crude intermediate was heated at 15 140 °C for 8 minutes to give a brown sticky oil that was then partly dissolved in dichloromethane. The solution was decanted from the mixture and concentrated to give the subtitle compound as an oil (673 mg).

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 8.61 (2H, s); 2.25 (1H, m); 1.13 (4H, m).

APCI-MS m/z: 199/201 1:1 [MH<sup>+</sup>].

20

b) (5S)-5-({[4-Ethynyl-3,6-dihydropyridin-1(2H)-yl]sulfonyl}methyl)-5-methylimidazolidine-2,4-dione

(5S)-5-Methyl-5-({[4-[(trimethylsilyl)ethynyl]-3,6-dihydropyridin-1(2H)-yl]sulfonyl}methyl)imidazolidine-2,4-dione (2.27 g, 6.0 mmol) and potassium fluoride 25 (1.07 g, 18.4 mmol) were stirred overnight at room temperature in methanol (50 mL). The solvent was evaporated off, the residue dissolved in EtOAc, washed with water followed by brine, dried (sodium sulphate) and evaporated. The residue was purified by column chromatography eluting with iso-hexane/EtOAc 1:1 to give a solid product (1.81 g).

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 1.66 (3H, s), 2.37 (2H, dt), 2.95 (1H, s), 3.24 - 3.50 (4H, m), 3.89 30 (2H, t), 6.11 (1H, s), 6.68 (1H, s), 8.75 (1H, s).

APCI-MS m/z: 298 [MH<sup>+</sup>].

c) (5S)-5-Methyl-5-({4-[(trimethylsilyl)ethynyl]-3,6-dihydropyridin-1(2H)-yl}sulfonyl)methylimidazolidine-2,4-dione

4-[(Trimethylsilyl)ethynyl]-1,2,3,6-tetrahydropyridine hydrochloride (3.43 g, 15.9 mmol) was stirred in THF (100 mL) with [(4S)-4-methyl-2,5-dioxoimidazolidin-4-yl]methanesulfonyl chloride (3.39 g, 15 mmol) and cooled in an ice salt bath (temperature about -10 °C). *N*-Ethyl-diisopropylamine (5.13 mL, 30 mmol) in THF (100 mL) was added dropwise over 2 hours and the mixture stirred a further 2 hours. The reaction mixture was washed with water, the aqueous layer extracted into EtOAc (×2), the organic phases combined, washed with 2M HCl (×2), saturated bicarbonate solution (×2), followed by brine, dried (sodium sulphate) and evaporated to give the crude product (5.06 g). This was used without further purification.

<sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 10.74 (1H, s), 8.01 (1H, s), 6.13 (1H, quintet), 3.75 (2H, d), 3.44 (2H, dd), 3.23 (2H, t), 2.18 - 2.28 (2H, m), 1.32 (3H, s), 1.32 (9H, s).  
APCI-MS *m/z*: 370 [MH<sup>+</sup>].

d) 4-[(Trimethylsilyl)ethynyl]-1,2,3,6-tetrahydropyridine hydrochloride

*tert*-Butyl 4-[(trimethylsilyl)ethynyl]-3,6-dihydropyridine-1(2H)-carboxylate (2.75 g, 9.8 mmol) was stirred in methanol (10 mL) and acetyl chloride (2.1 mL, 29.2 mmol) was added dropwise. The temperature rose from 18 °C to 30 °C during the addition, and the mixture was kept at 40 °C until there was no more starting material by tlc. The mixture was cooled to room temperature, EtOAc (15 mL) was added and the solid filtered off to give an off-white solid (1.6 g).

<sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 9.46 (2H, s), 6.09 (1H, quintet), 3.60 (2H, dd), 3.13 (2H, t), 2.35 (2H, td), 0.17 (8H, s).  
APCI-MS *m/z*: 180 [MH<sup>+</sup>].

e) *tert*-Butyl 4-[(trimethylsilyl)ethynyl]-3,6-dihydropyridine-1(2H)-carboxylate

Prepared from *N*-Boc-piperidin-4-one as in WO 96/05200.

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 6.05 (1H, s), 3.94 (2H, dd), 3.47 (2H, t), 2.23 (2H, dq), 1.45 (10H, s), 0.15 (8H, s).

GCMS-MS m/z: 223 [M-55].

f) [(4S)-4-Methyl-2,5-dioxoimidazolidin-4-yl]methanesulfonyl chloride

Prepared according to methods described in the following publications: Mosher, J., *J. Org. Chem.* **1958**, *23*, 1257; Griffith, O., *J. Biol. Chem.* **1983**, *258*, (3), 1591; and WO 02/074767.

Example 2

10 (5S)-5-Methyl-5-([4-{[2-(methylthio)pyrimidin-5-yl]ethynyl}-3,6-dihydropyridin-1(2H)-yl]sulfonyl)methyl)imidazolidine-2,4-dione

The title compound was prepared by the general method described by Nishihara et al., *J. Org. Chem.*, **2000**, *65*, 1780-1787. To a solution of 2-(methylthio)-5-  
15 [(trimethylsilyl)ethynyl]pyrimidine (0.55 g, 2.47 mmol) and 1-[(4-methyl-2,5-dioxoimidazolidin-4-yl)methyl]sulfonyl-1,2,3,6-tetrahydropyridin-4-yl trifluoromethanesulfonate (0.94 g, 2.22 mmol) in DMF (5 mL) was added CuI (10 mol %) and PdCl<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub> (5 mol %) and the mixture was heated at 85 °C for 6 hours.

The mixture was partitioned between EtOAc (20 mL) and water (10 mL), and the aqueous  
20 layer was extracted three times with EtOAc. The combined organic layers were washed with brine, water and concentrated to a brown oil (1.6 g). The title compound was obtained as a solid (10 mg) following purification by preparative HPLC (using a Xterra-Prep-MS-C18 (50 × 19) column with a 12 minutes gradient 5-35% of acetonitrile in water with 0.06% NH<sub>3</sub>).

25 <sup>1</sup>H-NMR (DMSO-*d*<sub>6</sub>): δ 10.75 (1H, s); 8.73 (2H, s); 8.02 (1H, s); 6.29 (1H, m); 3.84 (2H, m); 3.48 (2H, q); 3.30 (2H, m); 2.53 (3H, s); 2.38 (2H, m); 1.33 (3H, s).  
APCI-MS m/z: 422 [MH<sup>+</sup>].

a) 5-Bromo-2-(methylthio)pyrimidine

30 The subtitle compound was prepared following a method by Takahashi et al., *Chem. Pharm. Bull.*, **1958**, *6*, 334-337. To a solution of 5-bromo-2-chloropyrimidine (1.0 g, 5.2



mmol) in EtOH was added sodium methanethiolate (0.36 g, 5.2 mmol) at room temperature and the reaction mixture was stirred overnight. The mixture was partitioned between EtOAc (15 mL) and water (10 mL). The aqueous layer was extracted twice with EtOAc and washed with brine. The combined organic layers were dried and concentrated to give the subtitle compound as a white solid (1.1 g).

<sup>1</sup>H-NMR (CD<sub>3</sub>OD): δ 8.66 (2H, s); 2.54 (3H, s).

APCI-MS m/z: 204/206 1:1 [MH<sup>+</sup>].

b) 2-(Methylthio)-5-[(trimethylsilyl)ethynyl]pyrimidine

The subtitle compound was prepared following a method by Yamanaka et al, *Synth. Commun.*, **1983**, 312-314. To 5-bromo-2-(methylthio)pyrimidine (0.60 g, 2.9 mmol) in Et<sub>3</sub>N (3 mL) was added DMF (0.5 mL), CuI (5 mol %) and PdCl<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub> (3 mol %). The mixture was heated at 95 °C for 12 hours in a sealed tube and then partitioned between Et<sub>2</sub>O (30 mL) and water (10 mL). The aqueous layer was extracted twice with Et<sub>2</sub>O and the combined organic layers were washed with water, dried and concentrated to give the crude product as a brown oil. The compound was purified by flash chromatography using a gradient of 10-60% EtOAc in heptane, which gave the subtitle compound as a colourless oil (0.55 g).

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 8.56 (2H, s); 2.58 (3H, s); 0.27 (9H, s).

APCI-MS m/z: 223 [MH<sup>+</sup>].

c) 1-({[(4S)-4-Methyl-2,5-dioxoimidazolidin-4-yl]methyl}sulfonyl)-1,2,3,6-tetrahydropyridin-4-yl trifluoromethanesulfonate

4-({[(Trifluoromethyl)sulfonyl]oxy}-1,2,3,6-tetrahydropyridinium chloride was reacted with [(4S)-4-methyl-2,5-dioxoimidazolidin-4-yl]methanesulfonyl chloride (Example 1f) in the same way as for Example 1c.

<sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 10.77 (1H, s), 8.04 (1H, d), 6.10 (1H, t), 3.88 (2H, q), 3.36 - 3.58 (4H, m), 2.50 - 2.56 (2H, m), 1.32 (3H, s).

APCI-MS m/z: 422 [MH<sup>+</sup>].

d) 4-({[(Trifluoromethyl)sulfonyl]oxy}-1,2,3,6-tetrahydropyridinium chloride

*tert*-Butyl 4-[[[(trifluoromethyl)sulfonyl]oxy]-3,6-dihydropyridine-1(2*H*)-carboxylate (3.77 g, 11.4 mmol) was mixed with THF (15 mL) and concentrated hydrochloric acid (15 mL). After 1 hour, the mixture was evaporated and dried by azeotropic evaporation with toluene and methanol to give a beige solid (88 %) that could be used without further purification.

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 9.72 (2H, s), 6.22 (1H, s), 3.75 (2H, q), 3.30 (2H, t), 2.65 (2H, td).

APCI-MS m/z: 232 [MH<sup>+</sup>].

e) *tert*-Butyl 4-[[[(trifluoromethyl)sulfonyl]oxy]-3,6-dihydropyridine-1(2*H*)-carboxylate

A solution of *N*-*boc*-piperidin-4-one (10.14 g, 50 mmol) in THF (80 mL) was added to a cooled solution (-78 °C) of 2M LDA in THF (30 mL, 60 mmol, 1.2 eq.) and THF (80 mL) over approximately 30 minutes. After stirring a further 10 minutes, a solution of 1,1,1-trifluoro-*N*-phenyl-*N*-[(trifluoromethyl)sulfonyl]methanesulfonamide (20 g, 56 mmol, 1.1 eq.) in THF (80 mL) was added and the mixture was allowed to warm to room temperature. The solution was washed with water, the aqueous layer washed with EtOAc (×2), organic phases combined and washed with saturated ammonium chloride solution, brine, dried (sodium sulphate) and evaporated. The residue was filtered through neutral Alumina (200 g) eluting with *n*-heptane followed by *n*-heptane/EtOAc 9:1. After evaporation the <sup>1</sup>H-NMR showed some triflating agent still present but the crude product was used without further purification. Yield (13.17 g, 79.5%). (Wustrow, D. J., *Synthesis*, 1991, 993-995).

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 5.77 (1H, s), 4.05 (2H, q), 3.64 (2H, t), 2.45 (2H, quintet), 1.48 (9H, s).

GCMS-MS m/z: 274 [M-57].

**Example 3**

(5*S*)-5-Methyl-5-([4-{[2-(trifluoromethyl)pyrimidin-5-yl]ethynyl}-3,6-dihydropyridin-1(2*H*)-yl]sulfonyl)methylimidazolidine-2,4-dione

The title compound was prepared in 48% yield from 2-(trifluoromethyl)-5-[(trimethylsilyl)ethynyl]pyrimidine by the same method as described for Example 2. White solid from 95% EtOH, decomp. 240-245 °C.

<sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 10.8 (1H, br s), 9.16 (2H, s), 8.05 (1H, s), 6.42 (1H, m), 3.88 (2H, m), 3.56 (1H, d), 3.42 (1H, d), 3.32 (2H, m), 2.41 (2H, m) and 1.33 (3H, s).

APCI-MS m/z: 444 [MH<sup>+</sup>].

a) 2-(Trifluoromethyl)-5-[(trimethylsilyl)ethynyl]pyrimidine

2-(Trifluoromethyl)pyrimidine-5-yl trifluoromethanesulfonate (0.45 g, 1.5 mmol) and dry triethylamine (1.0 mL) were mixed in a screw-cap vial. The solution was purged with dry argon for 10 minutes. Trimethylsilylacetylene (0.43 mL, 3.0 mmol), finely ground CuI (0.010 g, 0.05 mmol) and PdCl<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub> (0.020 g, 0.030 mmol) were added. The vial was sealed and heated in an aluminum block at 80 °C. After stirring for 5 hours the volatiles were evaporated at room temperature (the product sublimes at 35-40 °C/ 10 mbar). The black residue was taken up in EtOAc (20 mL) and concentrated with silica (about 5 to 10 g) to dryness. Flash chromatography on silica with EtOAc/heptane (1:30) afforded 2-(trifluoromethyl)-5-[(trimethylsilyl)ethynyl]pyrimidine as a white solid (0.35 g, 95%), m.p. 75.5-76.0 °C.

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 8.90 (2H, s) and 0.30 (9H, s).

APCI-MS m/z: 245 [MH<sup>+</sup>].

b) 2-(Trifluoromethyl)pyrimidine-5-yl trifluoromethanesulfonate

Triflic anhydride (1.01 mL, 6.0 mmol) was added dropwise to a stirred mixture of 2-(trifluoromethyl)pyrimidin-5-ol (prepared according to US Patent 4,558,039) (0.82 g, 5.0 mmol), toluene (10 mL) and aqueous tripotassium phosphate (30% by weight, 10 mL) at ice-bath temperature (Frantz et al., *Organic Letters*, 2002, 4(26), 4717-4718). When the addition was complete the ice-bath was taken away and the solution was stirred at ambient temperature for 30 minutes. The clear phases were separated and the organic layer was washed with water, then brine. Drying of the organic phase over anhydrous sodium sulfate, filtration and concentration by rotary evaporation at room temperature afforded

2-(trifluoromethyl)-pyrimidine-5-yl trifluoromethanesulfonate as a colourless oil (1.38 g, 93%). B.p. 75-77 °C (10 mbar).

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 8.90 (2H, s).

5

#### Example 4

(5S)-5-Methyl-5-([4-[(2-methylpyrimidin-5-yl)ethynyl]-3,6-dihydropyridin-1(2H)-yl]sulfonyl)methylimidazolidine-2,4-dione

10 *tert*-Butyl 4-[(2-methylpyrimidin-5-yl)ethynyl]-3,6-dihydropyridine-1(2H)-carboxylate was treated with TFA in EtOH and after completion of the reaction the solvent was evaporated and the mixture was freeze dried. The residue was taken up in DMF (1.5 mL) and the mixture was cooled to 4 °C. *N*-Ethyl-diisopropylamine (2.2 eq.) was added and the mixture was stirred for 20 minutes before adding [(4S)-4-methyl-2,5-dioxoimidazolidin-4-yl]methanesulfonyl chloride (Example 1f) (1.1 eq.) in DMF (1 mL). The mixture was  
15 stirred for 10 min at 4 °C and then stirred for 2 h at room temperature before the solvent was evaporated. The product was purified by preparative HPLC to give the title compound (0.022 g, 30%).

<sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>); 10.75 (1H, s); 8.80 (2H, s); 8.02 (1H, s); 7.80 (1H, m); 7.32 (1H, d, *J* = 8.1 Hz); 6.24 (1H, s); 3.81 (2H, d, *J* = 3.2 Hz); 3.34 - 3.21 (2H, m); 3.30 (3H, s); 2.75 (2H, q, *J* = 20.8 Hz); 2.34 (2H, m); 1.29 (3H, s); 1.19 (3H, t, *J* = 7.6 Hz).  
20 APCI-MS *m/z*: 390 [MH<sup>+</sup>].

#### a) 2-Methyl-5-[(trimethylsilyl)ethynyl]pyrimidine

25 5-Bromo-2-methyl-pyrimidine (prepared according to UK patent application GB 2 157 288) (0.2 g, 1.16 mmol), (trimethylsilyl)acetylene (164 μL, 1.3 mmol), CuI (0.022 g, 0.116 mmol) and PdCl<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub> (0.082 g, 0.116 mmol) in Et<sub>3</sub>N (2 mL) and THF (2 mL) were stirred at 80 °C for 4 h. After cooling, the solvents were removed under vacuum and the residue chromatographed to give the subtitle compound (0.16 g, 50%).

30 APCI-MS *m/z*: 191 [MH<sup>+</sup>].

b) *tert*-Butyl 4-[(2-methylpyrimidin-5-yl)ethynyl]-3,6-dihydropyridine-1(2*H*)-carboxylate

To a solution of CuCl (1 mg, 0.01 mmol) and PdCl<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub> (0.003 g, 0.004 mmol) in DMF (2 mL) were added 2-methyl-5-[(trimethylsilyl)ethynyl]pyrimidine (0.088 g, 0.462 mmol) and *tert*-butyl 4-[[[(trifluoromethyl)sulfonyl]oxy]-3,6-dihydropyridine-1(2*H*)-carboxylate (Example 2e) (0.183 g, 0.555 mmol) at room temperature. The reaction mixture was stirred for 8 h at 80 °C. After cooling, the mixture was quenched with 1N HCl and extracted with ether (3 ×). The combined organic layers were washed with saturated aqueous NaHCO<sub>3</sub> solution, brine and dried. Filtration and evaporation afforded a brown oil, which was purified on HPLC to give the subtitle compound (0.062 g, 45%).  
APCI-MS m/z: 300 [MH<sup>+</sup>].

Example 5

15 (5*S*)-5-([4-[(2-Ethylpyrimidin-5-yl)ethynyl]-3,6-dihydropyridin-1(2*H*)-yl)sulfonyl}methyl)-5-methylimidazolidine-2,4-dione trifluoroacetate

The title compound was made in the same way as described for Example 4. The 5-bromo-2-ethyl-pyrimidine starting material was prepared using the methodology of GB 2 157 288.  
20 <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>); 10.75 (1H, s); 8.82 (2H, s); 8.05 (1H, s); 6.24 (1H, s); 3.81 (2H, d, *J* = 3.2 Hz); 3.34 - 3.21 (2H, m); 3.30 (3H, s); 2.92 (2H, q, *J* = 17.8 Hz); 2.75 (2H, q, *J* = 20.8 Hz); 2.34 (2H, m); 1.26 (3H, t, *J* = 12.8 Hz); 1.19 (1H, s).  
APCI-MS m/z: 404 [MH<sup>+</sup>].

25

Example 6

(5*S*)-5-([4-[(2-Methoxypyrimidin-5-yl)ethynyl]-3,6-dihydropyridin-1(2*H*)-yl)sulfonyl}methyl)-5-methylimidazolidine-2,4-dione

The title compound was prepared from 5-bromo-2-methoxypyrimidine and (5*S*)-5-{[(4-ethynyl-3,6-dihydropyridin-1(2*H*)-yl)sulfonyl]methyl}-5-methylimidazolidine-2,4-dione in the same way as described in Example 1.

5 <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>): δ 10.75 (1H, s); 8.73 (2H, s); 8.04 (1H, s); 6.26 (1H, s); 3.95 (3H, s); 3.81 (2H, d, *J* = 3.2 Hz); 3.34 - 3.21 (2H, m); 3.30 (3H, s); 2.75 (2H, q, *J* = 20.8 Hz); 2.34 (2H, m); 1.19 (3H, t, *J* = 7.6 Hz).  
APCI-MS *m/z*: 406 [MH<sup>+</sup>].

10

### Pharmacological Example

#### Isolated Enzyme Assays

##### 15 MMP12

Recombinant human MMP12 catalytic domain may be expressed and purified as described by Parkar A.A. *et al*, (2000), Protein Expression and Purification, 20, 152. The purified enzyme can be used to monitor inhibitors of activity as follows: MMP12 (50 ng/ml final concentration) is incubated for 60 minutes at room temperature with the synthetic substrate  
20 Mca-Pro-Cha-Gly-Nva-His-Ala-Dpa-NH<sub>2</sub> (10 μM) in assay buffer (0.1M "Tris-HCl" (trade mark) buffer, pH 7.3 containing 0.1M NaCl, 20mM CaCl<sub>2</sub>, 0.020 mM ZnCl and 0.05% (w/v) "Brij 35" (trade mark) detergent) in the presence (10 concentrations) or absence of inhibitors. Activity is determined by measuring the fluorescence at λ<sub>ex</sub> 320 nm and λ<sub>em</sub> 405 nm. Percent inhibition is calculated as follows:

25 % Inhibition is equal to the [Fluorescence<sub>plus inhibitor</sub> - Fluorescence<sub>background</sub>] divided by the [Fluorescence<sub>minus inhibitor</sub> - Fluorescence<sub>background</sub>].

##### MMP8

Purified pro-MMP8 is purchased from Calbiochem. The enzyme (at 10 μg/ml) is activated  
30 by p-amino-phenyl-mercuric acetate (APMA) at 1 mM for 2.5 h, 35 °C. The activated

enzyme can be used to monitor inhibitors of activity as follows: MMP8 (200 ng/ml final concentration) is incubated for 90 minutes at 35 °C (80% H<sub>2</sub>O) with the synthetic substrate Mca-Pro-Cha-Gly-Nva-His-Ala-Dpa-NH<sub>2</sub> (12.5 μM) in assay buffer (0.1M "Tris-HCl" (trade mark) buffer, pH 7.5 containing 0.1M NaCl, 30mM CaCl<sub>2</sub>, 0.040 mM ZnCl and 0.05% (w/v) "Brij 35" (trade mark) detergent) in the presence (10 concentrations) or absence of inhibitors. Activity is determined by measuring the fluorescence at λ<sub>ex</sub> 320 nm and λ<sub>em</sub> 405 nm. Percent inhibition is calculated as follows:

% Inhibition is equal to the  $[\text{Fluorescence}_{\text{plus inhibitor}} - \text{Fluorescence}_{\text{background}}]$  divided by the  $[\text{Fluorescence}_{\text{minus inhibitor}} - \text{Fluorescence}_{\text{background}}]$ .

#### MMP9

Recombinant human MMP9 catalytic domain was expressed and then purified by Zn chelate column chromatography followed by hydroxamate affinity column chromatography. The enzyme can be used to monitor inhibitors of activity as follows:

MMP9 (5 ng/ml final concentration) is incubated for 30 minutes at RT with the synthetic substrate Mca-Pro-Cha-Gly-Nva-His-Ala-Dpa-NH<sub>2</sub> (5 μM) in assay buffer (0.1M "Tris-HCl" (trade mark) buffer, pH 7.3 containing 0.1M NaCl, 20mM CaCl<sub>2</sub>, 0.020 mM ZnCl and 0.05% (w/v) "Brij 35" (trade mark) detergent) in the presence (10 concentrations) or absence of inhibitors. Activity is determined by measuring the fluorescence at λ<sub>ex</sub> 320 nm and λ<sub>em</sub> 405 nm. Percent inhibition is calculated as follows:

% Inhibition is equal to the  $[\text{Fluorescence}_{\text{plus inhibitor}} - \text{Fluorescence}_{\text{background}}]$  divided by the  $[\text{Fluorescence}_{\text{minus inhibitor}} - \text{Fluorescence}_{\text{background}}]$ .

#### MMP14

Recombinant human MMP14 catalytic domain may be expressed and purified as described by Parkar A.A. *et al*, (2000), Protein Expression and Purification, 20, 152. The purified enzyme can be used to monitor inhibitors of activity as follows: MMP14 (10 ng/ml final concentration) is incubated for 60 minutes at room temperature with the synthetic substrate Mca-Pro-Cha-Gly-Nva-His-Ala-Dpa-NH<sub>2</sub> (10 μM) in assay buffer (0.1M "Tris-HCl" (trade mark) buffer, pH 7.5 containing 0.1M NaCl, 20mM CaCl<sub>2</sub>, 0.020 mM ZnCl and

0.05% (w/v) "Brij 35" (trade mark) detergent) in the presence (5 concentrations) or absence of inhibitors. Activity is determined by measuring the fluorescence at  $\lambda_{ex}$  320 nm and  $\lambda_{em}$  405 nm. Percent inhibition is calculated as follows: % Inhibition is equal to the [Fluorescence<sub>plus inhibitor</sub> - Fluorescence<sub>background</sub>] divided by the [Fluorescence<sub>minus inhibitor</sub> - Fluorescence<sub>background</sub>].

A protocol for testing against other matrix metalloproteinases, including MMP9, using expressed and purified pro MMP is described, for instance, by C. Graham Knight *et al.*, (1992) FEBS Lett., 296(3), 263-266.

### MMP19

Recombinant human MMP19 catalytic domain may be expressed and purified as described by Parkar A.A. *et al.*, (2000), Protein Expression and Purification, 20:152. The purified enzyme can be used to monitor inhibitors of activity as follows: MMP19 (40 ng/ml final concentration) is incubated for 120 minutes at 35 °C with the synthetic substrate Mca-Pro-Leu-Ala-Nva-Dpa-Ala-Arg-NH<sub>2</sub> (5  $\mu$ M) in assay buffer (0.1M "Tris-HCl" (trade mark) buffer, pH 7.3 containing 0.1M NaCl, 20mM CaCl<sub>2</sub>, 0.020 mM ZnCl and 0.05% (w/v) "Brij 35" (trade mark) detergent) in the presence (5 concentrations) or absence of inhibitors. Activity is determined by measuring the fluorescence at  $\lambda_{ex}$  320 nm and  $\lambda_{em}$  405 nm. Percent inhibition is calculated as follows: % Inhibition is equal to the [Fluorescence<sub>plus inhibitor</sub> - Fluorescence<sub>background</sub>] divided by the [Fluorescence<sub>minus inhibitor</sub> - Fluorescence<sub>background</sub>].

### Protein Binding

Plasma protein binding was determined by ultrafiltration in an automated 96 well format assay. On each test occasion the plasma protein binding of a reference compound (budesonide) was monitored in parallel.

Test compounds (10 mM dissolved in DMSO) were added to plasma to a final concentration of 10  $\mu$ M and equilibrated at room temperature for 10 minutes. 350  $\mu$ L of



the plasma was transferred to an ultrafiltration plate, Microcon-96 (10kDa cutoff, Millipore). The ultrafiltration plate was centrifuged at 3000G for 70 minutes at room temperature. After centrifugation, the concentration of the compounds in the obtained plasma water (the unbound fraction) was determined by LC-MS/MS using a 3-point calibration curve and compared to the concentration in the original spiked plasma.

The analyses were performed using a gradient chromatographic system with acetic acid/acetonitrile as mobile phases. The detection was done using a triple quadrupole mass spectrometer, API3000 or API4000, from Applied Biosystems, with an electrospray interface.

#### **Protocol for Determination of Solubility**

The solubility of test compounds in 0.1M phosphate buffer, pH 7.4, was determined as follows:

The test compound (1 mg) was weighed into a 2 mL glass vial with a screw cap and 0.1M phosphate buffer pH 7.4. (1.00 mL) was added. The sample vial was then sonicated for about 10 minutes and then placed on a shake board overnight at 20 °C. The contents of the sample vial were then filtered through a Millipore Millex-LH 0.45 µm filter into a new 2 mL glass vial to give a clear solution. The clear solution (40 µL) was transferred to a new 2 mL glass vial and diluted with 0.1M phosphate buffer, pH 7.4 (960 µL).

A standard calibration curve for each particular test compound was established using solutions of known concentration. These solutions of known concentration were normally chosen to have concentrations of ~10 µg/mL and ~50 µg/mL. They were prepared by dissolving a known weight of the compound in 99.5 % ethanol (500 µL) and then sonicating for one minute if necessary. If the compound was still not completely dissolved, DMSO (500 µL) was added and the mixture sonicated for an additional one minute. The resulting solution was then diluted to the appropriate volume with a mixture of acetonitrile/100 mM ammonium acetate pH 5.5 20-50/80-50. If necessary, a further, more dilute, standard solution was prepared by dilution.

Test compound solutions and standard solutions were then analysed by HPLC with UV-detection using the following parameters and the solubility of the test compound in 0.1M phosphate buffer was thereby determined:

5

HPLC-equipment: HP1100/HP1050

Column: HyPURITY Advanced, 5  $\mu$ m, 125 x 3mm

Column temperature: RT

Flow rate: 1 mL/min

10

Mobile phase: A = acetonitrile

B = 100 mM ammonium acetate pH 5.5

Isocratic ratio: A/B 20-50/80-50

UV detector: 254 nm (220-280 nm)

Injection volume: 20  $\mu$ L

15

Chromatographic data handling system: ATLAS/Xchrome

#### **Protocol for Determination of Log D**

Log D values at pH 7.4 were determined using a shake flask method. An appropriate small  
20 amount of the test compound was placed in a 2 mL glass vial with a screw cap at room  
temperature and 600  $\mu$ L of 1-octanol (saturated with 10 mM phosphate buffer pH 7.4) was  
added. The vial was then sonicated for one minute so as to dissolve the compound  
completely. Then 600  $\mu$ L of 10 mM phosphate buffer pH 7.4 (saturated with 1-octanol)  
was added and the vial was shaken for 4 minutes to mix the two phases. The two phases  
25 were then separated by centrifugation of the sample at 1000g for 10 minutes at room  
temperature. Finally, the separated aqueous and organic phases were analysed in duplicate  
by HPLC using the following conditions:

Injector: Spark Holland, Endurance

Pump: HP1050

30

Detector: Kratos, Spectroflow 783

Column: YMC Pro C18, 5  $\mu$ m, 50x4mm, Part no. AS12S050504QT

Column temperature: RT

Flow rate:	1 mL/min
Mobile phase:	A = acetonitrile
	B = 25 mM formic acid
	C = 100 mM ammonium acetate pH 5.5
5	D = 0.05 % ammonium acetate
Gradient:	0.00 min    A/B or A/C or A/D    5/95
	5.00 min    A/B or A/C or A/D    100/0
	7.00 min    A/B or A/C or A/D    100/0
	7.02 min    A/B or A/C or A/D    5/95
10	UV detector: 254 nm
	Injection volume: 50 µL of undiluted aqueous phase and 5 µL of 10 times diluted (with methanol) organic phase
	Injection cycle time: 11 min
	Centrifuge: Hettich, Universal 30RF
15	Vortex: Scientific Industries, Vortex-2 genie
	Chromatographic data handling system: ATLAS/Xchrome

The log  $D_{pH7.4}$  value was automatically calculated (see equation below) by an Excel sheet after manual typing of compound peak area responses which were reported from the

20 ATLAS chromatographic data handling system.

Calculation of log  $D_{pH7.4}$  by equation:

$$25 \quad \text{Log } D = \left( \frac{[\text{Analyte}]_{\text{org}}}{[\text{Analyte}]_{\text{aq}}} \right) = \log \left( \frac{\text{Area}_{\text{org}} \times \text{Dilution factor}_{\text{org}}}{\text{Area}_{\text{aq}} \times \text{Dilution factor}_{\text{aq}} \times \frac{V_{\text{inj}}(\text{org})}{V_{\text{inj}}(\text{aq})}} \right)$$


The following table shows data for a representative selection of the compounds of the present invention and for selected compounds from WO 02/074767.

Table

Compound	hMMP12 IC <sub>50</sub> (nM)	hMMP9 IC <sub>50</sub> (nM)	hMMP8 IC <sub>50</sub> (nM)	hMMP14 IC <sub>50</sub> (nM)	hMMP19 IC <sub>50</sub> (nM)	Solubility pH 7.4 (μM)	Protein binding (% free)
Example 5	9	9	1,140	>10,000	>10,000	91	47
Example 3	6	5	547	>10,000	7,900	118	33
Example 1	6	5	2,530	>10,000	>10,000	396	18
WO 02/074767, page 120 (5 <i>S</i> )-5-({[4-[(4-chlorophenyl)ethynyl]-3,6-dihydropyridin-1(2 <i>H</i> )-yl]sulfonyl}-methyl)-5-methyl-imidazolidine-2,4-dione	2	9	180	4,300	3,980	49	1.75
WO 02/074767, page 120 (5 <i>S</i> )-5-methyl-5-({[4-[(4-methylphenyl)ethynyl]-3,6-dihydropyridin-1(2 <i>H</i> )-yl]sulfonyl}-methyl)imidazolidine-2,4-dione	3	34	384	4,430	1,970	72	2.25

## CLAIMS

1. A compound of formula (I) or a pharmaceutically acceptable salt thereof



5

wherein

$R^1$  represents C1 to 2 alkyl, cyclopropyl,  $OCH_3$ ,  $SCH_3$  or  $OCF_3$ ; said alkyl or cyclopropyl group being optionally further substituted by one or more fluoro atoms; and

10

$R^2$  represents C1 to 3 alkyl.

2. A compound according to claim 1, wherein  $R^1$  represents C1 to 2 alkyl or cyclopropyl; said alkyl or cyclopropyl group being optionally further substituted by one or more fluoro atoms.

15

3. A compound according to claim 2, wherein  $R^1$  represents C1 to 2 alkyl optionally further substituted by one or more fluoro atoms.

4. A compound according to Claim 3, wherein  $R^1$  represents  $CF_3$ .

20

5. A compound according to Claim 2, wherein  $R^1$  represents cyclopropyl.

6. A compound according to any one of claims 1 to 5, wherein  $R^2$  represents methyl or ethyl.

5 7. A compound according to Claim 6, wherein  $R^2$  represents methyl.

8. A compound according to claim 1 which is selected from the group consisting of:

(5*S*)-5-({[4-[(2-cyclopropylpyrimidin-5-yl)ethynyl]-3,6-dihydropyridin-1(2*H*)-yl]sulfonyl}methyl)-5-methylimidazolidine-2,4-dione;

10 (5*S*)-5-methyl-5-({[4-[[2-(methylthio)pyrimidin-5-yl]ethynyl]-3,6-dihydropyridin-1(2*H*)-yl]sulfonyl}methyl)imidazolidine-2,4-dione;

(5*S*)-5-methyl-5-({[4-[[2-(trifluoromethyl)pyrimidin-5-yl]ethynyl]-3,6-dihydropyridin-1(2*H*)-yl]sulfonyl}methyl)imidazolidine-2,4-dione;

15 (5*S*)-5-methyl-5-({[4-[(2-methylpyrimidin-5-yl)ethynyl]-3,6-dihydropyridin-1(2*H*)-yl]sulfonyl}methyl)imidazolidine-2,4-dione;


(5*S*)-5-({[4-[(2-ethylpyrimidin-5-yl)ethynyl]-3,6-dihydropyridin-1(2*H*)-yl]sulfonyl}methyl)-5-methylimidazolidine-2,4-dione;

(5*S*)-5-({[4-[(2-methoxypyrimidin-5-yl)ethynyl]-3,6-dihydropyridin-1(2*H*)-yl]sulfonyl}methyl)-5-methylimidazolidine-2,4-dione;

20 and pharmaceutically acceptable salts thereof.


9. A process for the preparation of a compound of formula (I) as defined in claim 1 or a pharmaceutically acceptable salt thereof which comprises:

a) reaction of a compound of formula (II)



(II)


wherein  $R^2$  is as defined in formula (I) and  $L^1$  represents a leaving group, with a compound of formula (III) (or a salt thereof)




5

wherein  $R^1$  is as defined in formula (I); or

b) reaction of a compound of formula (X)




10 wherein  $R^2$  is as defined in formula (I),  $R^3$  is H or a suitable protecting group and X is a leaving group such as halide or triflate; with an acetylenic compound of formula (IX)




15 wherein  $R^1$  is as defined in formula (I); or

c) reaction of a compound of formula (XI)

39



wherein R represents H or trimethylsilyl, R<sup>2</sup> is as defined in formula (I) and R<sup>3</sup> represents H or a suitable protecting group; with an aryl halide or triflate of formula (VI)



5

wherein R<sup>1</sup> is as defined in formula (I) and X represents halide or triflate;  
and optionally thereafter forming a pharmaceutically acceptable salt thereof.

10 10. A pharmaceutical composition comprising a compound of formula (I) or a pharmaceutically acceptable salt thereof as claimed in any one of claims 1 to 8 in association with a pharmaceutically acceptable adjuvant, diluent or carrier.

11. A process for the preparation of a pharmaceutical composition as claimed in claim 10  
15 which comprises mixing a compound of formula (I) or a pharmaceutically acceptable salt thereof as defined in any one of claims 1 to 8 with a pharmaceutically acceptable adjuvant, diluent or carrier.

12. A compound of formula (I) or a pharmaceutically acceptable salt thereof as claimed  
20 in any one of claims 1 to 8 for use in therapy.



13. Use of a compound of formula (I) or a pharmaceutically acceptable salt thereof as claimed in any one of claims 1 to 8 in the manufacture of a medicament for use in the treatment of an obstructive airways disease.

5

14. Use according to claim 13, wherein the obstructive airways disease is asthma or chronic obstructive pulmonary disease.

15. A method of treating a disease or condition mediated by MMP12 and/or MMP9 which comprises administering to a patient a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt thereof as claimed in any one of claims 1 to 8.

16. A method of treating an obstructive airways disease which comprises administering to a patient a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt thereof as claimed in any one of claims 1 to 8.

# INTERNATIONAL SEARCH REPORT

In application No.  
PCT/SE 2005/001092

<b>A. CLASSIFICATION OF SUBJECT MATTER</b>		
IPC7: C07D 401/14, A61K 31/506, A61P 11/00 According to International Patent Classification (IPC) or to both national classification and IPC		
<b>B. FIELDS SEARCHED</b>		
Minimum documentation searched (classification system followed by classification symbols)		
IPC7: C07D, A61K		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
SE,DK,FI,NO classes as above		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)		
EPO-INTERNAL, WPI DATA, PAJ, CHEM ABS DATA		
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 02074767 A1 (ASTRAZENECA AB), 26 Sept 2002 (26.09.2002)	1-14
	--	
A	WO 02074750 A1 (ASTRAZENECA AB), 26 Sept 2002 (26.09.2002)	1-14
	--	
A	WO 02096426 A1 (BRISTOL-MYERS SQUIBB COMPANY), 5 December 2002 (05.12.2002)	1-14
	--	
	-----	
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input checked="" type="checkbox"/> See patent family annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier application or patent but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family		
Date of the actual completion of the international search		Date of mailing of the international search report
27 Sept 2005		29-09-2005
Name and mailing address of the ISA/ Swedish Patent Office Box 5055, S-102 42 STOCKHOLM Facsimile No. +46 8 666 02 86		Authorized officer EVA JOHANSSON/Els Telephone No. +46 8 782 25 00

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/SE2005/001092

## Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 15-16  
because they relate to subject matter not required to be searched by this Authority, namely:  
Claims 15-16 relate to a method of treatment of the human or animal body by surgery or by therapy, as well as diagnostic  
.../...
2. ☐ Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

### Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- ☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- ☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/SE2005/001092

Box II.1

methods /Rule 39.1(iv). Nevertheless, a search has been executed for these claims. The search has been based on the alleged effects of the compounds.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

31/08/2005

International application No.

PCT/SE 2005/001092

WO	02074767	A1	26/09/2002	AU	5282101 A	26/11/2001
				BR	0207983 A	15/06/2004
				BR	0207984 A	15/06/2004
				BR	0208104 A	02/03/2004
				BR	0208105 A	09/03/2004
				CA	2440473 A	26/09/2002
				CA	2440630 A	26/09/2002
				CA	2440631 A	26/09/2002
				CA	2440632 A	26/09/2002
				CN	1509272 A	30/06/2004
				CN	1509275 A	30/06/2004
				CN	1509276 A	30/06/2004
				CN	1509286 A	30/06/2004
				CZ	20032497 A	18/02/2004
				CZ	20032499 A	17/03/2004
				CZ	20032500 A	18/02/2004
				CZ	20032502 A	14/01/2004
				EE	200300439 A	15/12/2003
				EE	200300445 A	15/12/2003
				EE	200300449 A	15/12/2003
				EE	200300451 A	15/12/2003
				EP	1370534 A	17/12/2003
				EP	1370536 A	17/12/2003
				EP	1370537 A	17/12/2003
				EP	1370556 A	17/12/2003
				HU	0400194 A	28/07/2004
				HU	0400202 A	30/08/2004
				HU	0400206 A	30/08/2004
				HU	0400327 A	28/01/2005
				IL	157570 D	00/00/0000
				IL	157652 D	00/00/0000
				IL	157656 D	00/00/0000
				IL	157657 D	00/00/0000
				JP	2004523581 T	05/08/2004
				JP	2004523583 T	05/08/2004
				JP	2004527511 T	09/09/2004
				JP	2004527515 T	09/09/2004
				MX	PA03008177 A	12/12/2003
				MX	PA03008180 A	12/12/2003
				MX	PA03008181 A	12/12/2003
				MX	PA03008191 A	29/01/2004
				NO	20034025 A	13/11/2003
				NO	20034042 A	10/11/2003
				NO	20034044 A	10/11/2003
				NO	20034045 A	10/11/2003
				NZ	528106 A	24/03/2005
				NZ	528107 A	24/06/2005
				NZ	528140 A	25/02/2005
				PL	364706 A	13/12/2004
				PL	364707 A	13/12/2004
				PL	364714 A	13/12/2004
				PL	365099 A	27/12/2004
				SE	0100902 D	00/00/0000
				SK	10922003 A	04/05/2004
				SK	10932003 A	06/04/2004

# INTERNATIONAL SEARCH REPORT

Information on patent family members

31/08/2005

International application No.

PCT/SE 2005/001092

WO	02074767	A1	26/09/2002	SK	10952003	A	04/05/2004
				SK	10962003	A	02/03/2004
				US	20040106659	A	03/06/2004
				US	20040127528	A	01/07/2004
				US	20040138276	A	15/07/2004
				US	20040147573	A	29/07/2004
				WO	02074748	A	26/09/2002
				WO	02074750	A	26/09/2002
				WO	02074751	A	26/09/2002
				ZA	200306731	A	29/11/2004
				ZA	200306732	A	29/11/2004
				ZA	200306734	A	29/11/2004
				ZA	200306737	A	29/11/2004

---

## INTERNATIONAL SEARCH REPORT

Information on patent family members

31/08/2005

International application No.

PCT/SE 2005/001092

WO	02074750	A1	26/09/2002	AU	5282101	A	26/11/2001
				BR	0207983	A	15/06/2004
				BR	0207984	A	15/06/2004
				BR	0208104	A	02/03/2004
				BR	0208105	A	09/03/2004
				CA	2440473	A	26/09/2002
				CA	2440630	A	26/09/2002
				CA	2440631	A	26/09/2002
				CA	2440632	A	26/09/2002
				CN	1509272	A	30/06/2004
				CN	1509275	A	30/06/2004
				CN	1509276	A	30/06/2004
				CN	1509286	A	30/06/2004
				CZ	20032497	A	18/02/2004
				CZ	20032499	A	17/03/2004
				CZ	20032500	A	18/02/2004
				CZ	20032502	A	14/01/2004
				EE	200300439	A	15/12/2003
				EE	200300445	A	15/12/2003
				EE	200300449	A	15/12/2003
				EE	200300451	A	15/12/2003
				EP	1370534	A	17/12/2003
				EP	1370536	A	17/12/2003
				EP	1370537	A	17/12/2003
				EP	1370556	A	17/12/2003
				HU	0400194	A	28/07/2004
				HU	0400202	A	30/08/2004
				HU	0400206	A	30/08/2004
				HU	0400327	A	28/01/2005
				IL	157570	D	00/00/0000
				IL	157652	D	00/00/0000
				IL	157656	D	00/00/0000
				IL	157657	D	00/00/0000
				JP	2004523581	T	05/08/2004
				JP	2004523583	T	05/08/2004
				JP	2004527511	T	09/09/2004
				JP	2004527515	T	09/09/2004
				MX	PA03008177	A	12/12/2003
				MX	PA03008180	A	12/12/2003
				MX	PA03008181	A	12/12/2003
				MX	PA03008191	A	29/01/2004
				NO	20034025	A	13/11/2003
				NO	20034042	A	10/11/2003
				NO	20034044	A	10/11/2003
				NO	20034045	A	10/11/2003
				NZ	528106	A	24/03/2005
				NZ	528107	A	24/06/2005
				NZ	528140	A	25/02/2005
				PL	364706	A	13/12/2004
				PL	364707	A	13/12/2004
				PL	364714	A	13/12/2004
				PL	365099	A	27/12/2004
				SE	0100902	D	00/00/0000
				SK	10922003	A	04/05/2004
				SK	10932003	A	06/04/2004

# INTERNATIONAL SEARCH REPORT

Information on patent family members

31/08/2005

I  
ial application No.  
PCT/SE 2005/001092

WO	02074750	A1	26/09/2002	SK	10952003	A	04/05/2004
				SK	10962003	A	02/03/2004
				US	20040106659	A	03/06/2004
				US	20040127528	A	01/07/2004
				US	20040138276	A	15/07/2004
				US	20040147573	A	29/07/2004
				WO	02074748	A	26/09/2002
				WO	02074751	A	26/09/2002
				WO	02074767	A	26/09/2002
				ZA	200306731	A	29/11/2004
				ZA	200306732	A	29/11/2004
				ZA	200306734	A	29/11/2004
				ZA	200306737	A	29/11/2004
				AU	5282201	A	12/11/2001
				BR	0207985	A	15/06/2004
				BR	0208062	A	02/03/2004
				CA	2440475	A	26/09/2002
				CA	2444526	A	26/09/2002
				CN	1509273	A	30/06/2004
				CN	1509274	A	30/06/2004
				CZ	20032498	A	17/03/2004
				CZ	20032501	A	18/02/2004
				EE	200300450	A	15/12/2003
				EE	200300452	A	16/02/2004
				EP	1370535	A	17/12/2003
				EP	1370538	A	17/12/2003
				HU	0400193	A	28/07/2004
				HU	0400328	A	28/09/2004
				IL	157650	D	00/00/0000
				IL	157658	D	00/00/0000
				JP	2004523582	T	05/08/2004
				JP	2004527512	T	09/09/2004
				MX	PA03008183	A	12/12/2003
				MX	PA03008187	A	29/01/2004
				NO	20034027	A	05/11/2003
				NO	20034032	A	10/11/2003
				NZ	528108	A	29/04/2005
				NZ	528141	A	27/05/2005
				PL	364705	A	13/12/2004
				PL	365107	A	27/12/2004
				SE	0100903	D	00/00/0000
				SK	10912003	A	04/05/2004
				SK	10942003	A	06/04/2004
				US	20040110809	A	10/06/2004
				US	20040116486	A	17/06/2004
				WO	02074749	A	26/09/2002
				WO	02074752	A	26/09/2002
				ZA	200306733	A	29/11/2004
				ZA	200306738	A	29/11/2004



# INTERNATIONAL SEARCH REPORT

Information on patent family members

31/08/2005

International application No.

PCT/SE 2005/001092

WO	02096426	A1	05/12/2002	CA	2447475	A	05/12/2002
				EP	1397137	A	17/03/2004
				JP	2004535411	T	25/11/2004
				US	6890915	B	10/05/2005
				US	6906053	B	14/06/2005
				US	20030130273	A	10/07/2003
				US	20040209874	A	21/10/2004
				US	20050171096	A	04/08/2005

---

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**